

## Phytochemicals in Cereals and their Potential Health Benefits-A Review

Remya Mohanraj

Houston Community College, Houston, Texas, USA

### \*Corresponding author

Remya Mohanraj, Houston Community College, Houston, Texas, USA

Submitted: 12 March 2020; Accepted: 10 Jun 2020; Published: 01 Sept 2020

### Abstract

Edible seeds of Poaceae are referred to as cereals. A lot of cereals have been the staple food for human populations from time immemorial and have been a significant source of energy and nutrients. Their consumption has always been linked to a wide array of health benefits that could be attributed to the rich amount of phytochemicals harbored in them. This review attempts to present a summary of the various phytochemicals present in cereals and their potential health benefits. This review has been compiled using information from various sources including scientific publications. Citations have been provided to credit the sources.

### Introduction

Cereals belong to the family Poaceae and are cultivated for their nutritionally beneficial seeds referred to as grains that are used for human consumption directly as well as indirectly via livestock feed [1, 2]. The type of cereal consumed varies according to geographical regions. They are consumed either as raw grains or as components of food substances and whole grain cereals are abundant in vitamins, minerals, carbohydrates, fats, oils and protein [3].

Cereal grains consumption contributes to greater than 50% of global everyday caloric intake [4]. Wheat, rice, maize and oats occupy top position in US diet. Whole grains such as these abound in nutrients and phytochemicals that promote health [5]. Figure 1 shows a cereal plant. Grain is derived following fertilization of the flowers in the inflorescence

This article aims to provide a comprehensive review on phytochemicals in cereals and their potential health benefits. Information presented here has been compiled from existing published articles.


### Composition of Whole Grain Cereals:

**The different layers of grain are shown in Figure 2.**

The endosperm being the largest portion contains carbohydrates, proteins, vitamins, and minerals. The germ has vitamins, some protein, minerals, and fats. The outer most layer, bran protects the other two layers and contains phenolic compounds, vitamins, and minerals [6].



**Figure 1:** A cereal plant from our garden showing the formation of grains



**Figure 2:** Grain - Structure

When all three layers of the grain are present, it is referred to as the whole grain and is copious in essential vitamins, minerals and phytochemicals. The process of refining removes most of the bran and some of the germ leading to loss of dietary fiber and other important compounds from whole grains. Therefore, in comparison to whole grains, refined grains contain a higher starch content [3, 7, 8]. Whole grains contain high concentrations of B vitamins, minerals, tocol, arginine, lysine and a variety of phytochemicals including phytates, phenolic compounds, avenanthramides, avenalumic acid that confer them with high antioxidant potential [8,90].

Cereals are able to exercise varying metabolic effects because of the presence of both soluble and insoluble fibres [9]. Barley, oats, apples, pears and most legumes are the major sources of soluble fibre [91]. In a review of clinical trials related to health benefits of cereals, Smith and Tucker throw light on how intervention trials have revealed improved glucose and lipid metabolism with soluble fibre [9]. Borneo and Leon, in their review on functional components of cereals, list dietary fiber, inulin, beta-glucan, resistant starch, carotenoids, phenolics, tocotrienols, and tocopherols as the most predominant phytochemicals that are responsible for the health enhancing properties of cereals [10]. Cereals are also a good source of proteins. When they are consumed along with legumes, there is a complementation of amino acids resulting in higher protein quality. There is a variation in protein content among cereals [11]. Maize, rice, and wheat are quite popular [12]. According to Morrison, starches from cereal endosperm may be unique due to elevated levels of monoacyl lipids within the starch granules [13].

### Health Benefits

Health benefits from cereals are attributed to the presence of various vitamins, minerals and phytochemicals in them. According to Hollaender et al, a diet rich in whole grains lowers LDL cholesterol and total cholesterol. They also point out that the most effective whole grain for lowering cholesterol is whole-grain oat [14]. Studies carried out by Heather et al demonstrate the effect of whole grain diet in lowering total, LDL and HDL cholesterol. They also reported that

a diet rich in whole grains decreased cardiovascular risk factors [15]. Giglio et al. conducted studies to demonstrate the cardio protective functions of polyphenols [16]. The presence of polyphenols in cereals has been well documented [17, 18, 38]. Whole grains help lower blood pressure, which in turn is an important risk factor for heart disease. A lower risk of hypertension among males who had a higher intake of whole grain breakfast cereal has been reported by Kochar et al [19].

In a recent review on the effect of intact cereal grain fiber on gut microbiota composition, Angie and Katie reported increased microbiota diversity as a consequence of intact cereal fiber consumption [20]. The authors recommend increasing cereal fiber consumption for general good health and for promoting gut microbiota diversity. Also, the beneficial effects of dietary fibers in bowel movements are quite well known.

Diets rich in whole grains are correlated with a 20-30% reduction of risk in the onset of type-2 diabetes. This is due to the presence of dietary fiber, vitamins, minerals and phytochemicals in wholegrains. A majority of phytochemicals act as antioxidants *in vitro* and alleviate oxidative stress and inflammation that are involved in the pathogenesis of type-2 diabetes [21].

Studies indicate that consumption of whole grains has a correlation to reduced obesity risk and weight gain [22]. Studies have also pointed out that an increased intake of dietary carbohydrates through ready-to-eat cereals aids in weight reduction [23, 24]. The beneficial effect of whole grains on body weight can be attributed to their higher fiber content [25]. Eating whole grains can help cut down on the amount of body fat and decrease waist circumference [15].

The role of cereals in combating cancer is being researched. These properties are attributed to the fibers and germ of the cereal [26, 27]. According to Makarem et al, whole grains and cereal fiber might have a protective effect against gastrointestinal cancers [28].

Ferulic acid present in whole-wheat grain or wheat bran may be responsible for its anti-inflammatory properties [29]. Small particle size wheat bran fraction may help manage inflammatory disorders outside the gut [30]. Lopes et al. observed that a combination of whole sorghum breakfast cereal and unfermented probiotic milk decreased the serum levels of C-reactive protein and malondialdehyde and increased superoxide dismutase and total antioxidant potential in individuals with chronic kidney disease [31]. Whole grains can contribute to an increased life span. Numerous studies have reported the correlation between increased intake of whole grains and reduced risk of mortality [32-35; 92].

### Phytochemistry of Cereals

The wide variety of phytochemicals harbored in cereals give credence to the array health benefits conferred by them. Prominent phytochemical compounds in whole-grain cereals include phenolic compounds, carotenoids, phytic acid, phytosterols, tocols, gamma-oryzanol, beta-glucan, vitamins and minerals [5,8, 37, 90, 96 ].

**Phenolic Compounds:** Phenolic compounds are a group of secondary metabolites in plants including flavonoids, phenolic acids, tannins, lignans, and coumarins [36]. Phenolic compounds in cereals include Phenolic acid, Flavonoids, Avenanthramides, Lignans, and

Alkylresorcinols [37]. A review by Rao et al indicated that whole grain pigmented cereals contained anthocyanins, proanthocyanidins, and their derivatives which conferred them a higher antioxidant potential. Geographical locations and variations in levels of salinity, water, and temperature also influenced the phenolic content [17]. Pasha et al. assessed the antioxidant activity and analysed phenolic acids in sorghum and millet. Total phenolic content was in the range of 0.166-0.362 mg gallic acid equivalent and 0.275-0.305 mg in sorghum and millet respectively. Chlorogenic acid was the major phenolic acid in sorghum while it was m-coumaric acid in millet. The results showed that sorghum possessed higher antioxidant activities and phenolic acids as compared to millet [39].

Rao et al. characterized polyphenols and assessed the antioxidant potential of seven barley varieties from Australia. It was observed that Prodelphinidin B3 possessed the most antioxidant activity. Other polyphenols that exhibited antioxidant activity included procyanidin, glycosides of catechin and flavan-3-ols. The results of the study emphasized the merits of consuming whole grain barley [38].

It was revealed that integrating classical chemotherapy with nutrients and polyphenols from dietary sources might increase potency and decrease adverse side effects of the antineoplastic drug [40]. Saharan et al. observed that rice and wheat had the highest levels of polyphenols and antioxidants followed by sorghum, oat and maize because of fermentation. They also demonstrated the antioxidant activity was related to total phenol and flavonoid content [18]. Cereals are abundant in free phenolic acids like ferulic, sinapic, p-coumaric, 2-hydroxycinnamic, 2, 4-dihydroxybenzoic, caffeic, syringic and vanillic acid [41,42,93]. It has been reported that soluble phenolic acids are responsible for the flavor of cereal foods [43].

Smeds et al. analyzed lignins from cereals and reported the dominant presence of 7-Hydroxymatairesinol, in wheat, triticale, oat, barley, millet, corn bran, and amaranth whole grain. They also reported the occurrence of Syringaresinol, a cereal lignin [44].

Other lignans like buddlenol derivatives and hedyotisol lignans have been reported in rye bran [45]. Lignans are abundant in the bran fraction [43, 46, 47]. In the order of total lignin content cereals have been listed as rye, wheat, triticale, oat, spelt wheat, wild rice, barley, quinoa, red rice, brown rice by Smeds et al [44].

Stasiuk and Kozubek presented a review including the antimicrobial and antiparasitic activity of Alkylresorcinols (phenolic lipids) [48]. Activity exhibited by barley seeds against *Aspergillus niger* and *Penicillium crysogenum* has been attributed to the presence of alkylresorcinols in them [49]. It has also been reported that alkylresorcinols from rye bran inhibits the growth of *Penicillium expansum* and *Neofabraea perennans* [50, 51].

Flavonoids are a category of phenolic compounds and they could potentially halt the progression of certain chronic diseases [52]. Major cereal crops produce biologically active and stable to hydrolyze flavone-C-glycosides [53]. Zieliński et al. and Michalska et al. reported the amount of total flavonoid content in wholegrain rye [54,55]. According to Ivanišova et al. the flavonoid content in coarse and fine rye bran fractions was 1910 and 2390 mg/kg quercetin equivalents respectively [56]. Michalska et al. reported that there is a 46 % distribution of flavonoids in bran [55]. Evidences point out

that dietary flavonoids contribute to the reduction of various diseases [57, 58]. Studies in oat groats by Peterson et al. demonstrated that avenanthramides are evenly distributed in the outer groat mass and their concentration is much lower in the starchy endosperm [65].

**Benzoxazinoid:** Katina et al. reported the presence of benzoxazinoid in rye grains and Pedersen et al. presented a detailed study on the content of benzoxazinoids in rye flours and rye bread [59, 60]. Quantitative data on benzoxazinoid content in wheat and rye fractions was published by Tanwir et al, [61]. They reported that the benzoxazinoids were concentrated in rye bran as DIBOA-diglucoside. Aneugenic effects of benzoxazinoids on human-derived liver cells was reported by Buchmann et al, [62]. Adhikari et al. presented a review on the presumptive health effects of benzoxazinoids and their conversion products. These compounds were reported to possess anti-inflammatory, anticancer, antimicrobial, and stimulatory effects on central nervous system and reproductive system [63]. Steffensen et al. presented the effect of benzoxazinoids from rye in prostrate cancer patients [64].

**Carotenoids:** Ndolo and Beta attempted to compare the distribution of carotenoids in endosperm, germ and aleurone fractions in different types of grains [66]. A significant variation in the concentrations of carotenoids was observed even though the cereal carotenoid composition was similar. In germ fractions, concentration of lutein and zeaxanthin were higher in non-corn cereals while in the endosperms, lutein and zeaxanthin contents were lower. In the aleurone layer, zeaxanthin levels were higher than lutein [66]. El sayed evaluated carotenoid composition and studied the effect of genotype and environment on lutein utilization on wheat species. It was observed that *Triticum monococcum* possessed high levels of all-trans-lutein [67].

**Phytic Acid:** Phosphorous in grains and oil seeds is stored in the form of phytic acid or inositol hexaphosphate [94]. 50-80% of the total phosphorus in various cereals is accounted for by phytic acid and is incorporated in the seed during ripening [68]. Phytic acid is capable of chelating ions like zinc, calcium and iron leading to low bioavailability of minerals [95].

**Phytosterols:** Phytosterols are a class of sterols mostly present in cell walls and membranes of plant. Earlier studies have reported that phytosterols from diet can reduce intestinal cholesterol absorption there by lowering serum LDL and total cholesterol concentrations [69-73]. Analysis of the sterol profiles of extracted lipids from rice bran, wheat bran, wheat germ, durum wheat, oat bran, oat hull, and corn fine fiber by Jiang and Wang showed that rice bran contained the maximum amount of lipids while the least amount of lipids were found in corn fine fiber. The dominant phytosterols were sitosterol, campesterol, and stigmasterol. Brassicasterol was found only in wheat samples, while appreciable quantities of cycloartenol and 24-methylenecycloartanol was found in rice bran oil [74].

**Tocols:** Tocols are naturally occurring antioxidants in plant based food including cereal grains [75]. Tocols include tocopherols and tocotrienols [76]. These compounds are present in noticeable quantities in most cereal grains including barley, oats, wheat, rye, rice [77-79].

**Beta-glucan:**  $\beta$ -glucan consists of D-glucose monomers linked with

glycosidic bonds at  $\beta$  (1-3), (1-4),  $\beta$  (1-6) and is a major component of endosperm and aleurone cell walls of oat, barley, rye and wheat [80-82]. Havrlentová and Kraic reported that barley and oat had the highest content of  $\beta$ -D-glucan and are suitable sources of  $\beta$ -D-glucan that is beneficial for health [83]. The nutritional value of  $\beta$ -glucan is influenced by its tendency to be water soluble by virtue of its irregular structure. Soluble fiber confers protection against cardiovascular diseases [80].  $\beta$ -glucan from cereals is capable of lowering LDL, total cholesterol and serum triglyceride levels [84].

**Minerals and vitamins:** In a recent study on polish population; Laskowski et al reported that cereals and cereal products contributed 64.1% manganese to the average Polish diet. This was followed by iron 34.1% and copper 31.3%; 20–30% zinc and phosphorus; 10–20% sodium, potassium and calcium. They also supplied 33.6% folate, 20-30% thiamin, 10-20% riboflavin, niacin and vitamin B6 [85].

**Aminoacids and Proteins:** Cereals abound in their aminoacids and proteins content. Excepting oats and rice, prolamins are the major endosperm storage proteins of all cereal grains [86, 87]. Majority of storage proteins in rice are glutelins and rice protein has elevated lysine levels. Sorghum contains mainly prolamins and gluten proteins in wheat confer the unique properties of dough formation. Oats are high in glutamine and asparagine and are mildly insufficient in cysteine and methionine [88]. Oats accumulate significant amounts of globulins and prolamins. A higher accumulation of globulins increases their protein quality [89].

### Conclusion

From time immemorial, the human diet has been comprised of cereals. These magic grains have been our source of energy and have provided us with the necessary nutrients. Consuming cereals in the right quantities and in the right form could help alleviate numerous health issues. Research focusing on deducing the properties of phytochemicals from whole grains is imperative for the formulation of cereal-based products that can potentially promote health of human populations.

### References

- Bender DA, Bender AE (1999) Bender's Dictionary of Nutrition and Food technology, 7th edition. Woodhead Publishing, Abington.
- BNF (British Nutrition Foundation) (1994) Starchy Foods in the Diet. BNF, London.
- Sarwar MH, Sarwar MF, Sarwar M, Niaz Ahmad Qadri, Safia Moghal (2013) The importance of cereals (Poaceae: Gramineae) nutrition in human health: A review. *Journal of Cereals and Oilseeds* 4: 32-35.
- Awika JM (2011). Major Cereal Grains Production and Use around the World. In: *Advances in Cereal Science: Implications to Food Processing and Health Promotion* (Editors: J.M. Awika, V. Piironen, S. Bean) 1089: 1-13.
- Slavin J (2003) why whole grains are protective: biological mechanisms. *Proc Nutr Soc* 62: 129-134.
- Okarter N, Liu RH (2010) Health benefits of whole grain phytochemicals. *Crit Rev Food Sci Nutr* 50: 193-208.
- Slavin J, Margaret C Martini, David R Jacobs, Jr, Len Marquart (1999) Plausible mechanisms for the protectiveness of whole grains. *The American Journal of Clinical Nutrition* 70: 459s-463s.
- Slavin J (2004) Whole grains and human health. *Nutrition Research Reviews* 17: 99-110.
- Smith CE, Tucker KL (2011) Health benefits of cereal fibre: a review of clinical trials. *Nutrition research reviews* 24: 118-131.
- Borneo R, León AE (2012) Whole grain cereals: functional components and health benefits. *Food Funct* 3: 110-119.
- Hackler LR (1985) Cereal Proteins in Human Nutrition. In: Lásztity R., Hidvégi M. (eds) *Amino Acid Composition and Biological Value of Cereal Proteins*. Springer, Dordrecht.
- Guerrieri N, Cavaletto M (2018) Cereals Proteins. In: Yada RY. (eds) *Proteins in food processing* (Second Edition). Woodhead Publishing Series in Food Science, Technology and Nutrition Elsevier, UK.
- Morrison WR (1988) Lipids in cereal starches: A review. *Journal of Cereal Science* 8: 1-15.
- Hollænder PL, Ross AB, Kristensen M (2015). Whole-grain and blood lipid changes in apparently healthy adults: a systematic review and meta-analysis of randomized controlled studies 3: 556-572.
- Heather I Katcher, Richard S Legro, Allen R Kunselman, Peter J Gillies, Laurence M Demers, et al. (2008) The effects of a whole grain-enriched hypocaloric diet on cardiovascular disease risk factors in men and women with metabolic syndrome. *The American Journal of Clinical Nutrition* 87: 79-90.
- Giglio RV, Patti AM, Cicero AFG, Lippi G, Rizzo M, et al. (2018) Polyphenols: Potential use in the prevention and treatment of cardiovascular diseases. *Current Pharmaceutical Design* 24: 239-258.
- Rao S, Schwarz LJ, Santhakumar AB, Chinkwo KA, Blanchard CL (2018) Cereal phenolic contents as affected by variety and environment (Review). *Cereal Chemistry* 95: 589-602.
- Saharan P, Sath PK, Singh Duhan J (2017) Comparative assessment of effect of fermentation on phenolics, flavanoids and free radical scavenging activity of commonly used cereals. *Biocatalysis and Agricultural Biotechnology* 12: 236-240.
- Kochar J, Gaziano JM, Djoussé L (2012) Breakfast cereals and risk of hypertension in the Physicians' Health Study I. *Clinical Nutrition* 31: 89-92.

- 
20. Angie Jefferson, Katie Adolphus (2019) The Effects of Intact Cereal Grain Fibers, Including Wheat Bran on the Gut Microbiota Composition of Healthy Adults: A Systematic Review. *Front. Nutr* 6: 33.
  21. Belobrajdic DP, Bird AR (2013) The potential role of phytochemicals in wholegrain cereals for the prevention of type-2 diabetes. *Nutr J* 12: 62.
  22. Harland JI, Garton LE (2008) Whole-grain intake as a marker of healthy body weight and adiposity. *Public Health Nutr* 11: 554-563.
  23. Kirk TR, Burkill S, Cursiter M (1997) Dietary fat reduction achieved by increasing consumption of a starchy food – an intervention study. *Eur J Clin Nutr* 51: 455-461.
  24. Crombie N, Kirk T (1999) Prevention of weight gain and blood cholesterol reduction after consumption of a high carbohydrate food in men. *Intl J Obesity* 23: 557.
  25. Howarth NC, Saltzman E, Roberts SB (2001) Dietary fiber and weight regulation. *Nutr Rev* 59: 129-139.
  26. Challier B, Perarnau JM, Viel JF (1998) Garlic, onion and cereal fibre as protective factors for breast cancer: A French case-control study. *Eur J Epidemiol* 14: 737.
  27. Matti Hakama E, Saxén A (1967) Cereal consumption and gastric cancer. *International journal of cancer* 2: 265-268.
  28. Makarem N, Nicholson JM, Bandera EV, McKeown NM, Parekh N (2016) Consumption of whole grains and cereal fiber in relation to cancer risk: A systematic review of longitudinal studies. *Nutr Rev* 74: 353-373.
  29. Anson NM, Selinheimo E, Havenaar R, Anna-Marja A, Ismo Mattila, et al. (2009) Bioprocessing of Wheat Bran Improves in vitro Bioaccessibility and Colonic Metabolism of Phenolic Compounds. *J Agric Food Chem* 57: 6148-6155.
  30. Surianoa F, Audrey MN, Joran V, Marta O, SophieLeclercq, TomVan de Wiele, Christophe MC, Patrice DC, Laure BB, Nathalie MD (2018) Particle size determines the anti-inflammatory effect of wheat bran in a model of fructose over-consumption: Implication of the gut microbiota *J Funct Foods* 41:155-162.
  31. Lopes RDCSO, Lima SLS, da Silva BP, Toledo RCL, Moreira MEDC, et al. (2018) Evaluation of the health benefits of consumption of extruded tannin sorghum with unfermented probiotic milk in individuals with chronic kidney disease *Food Research International* 107: 629-638.
  32. Wu H, Flint AJ, Qi Q, van Dam RM, Sampson LA, et al. (2005) Association between dietary whole grain intake and risk of mortality: two large prospective studies in US men and women. *JAMA Intern Med* 175: 373-384.
  33. Ma X, Tang WG, Yang Y, Zhang QL, Zheng JL, et al. (2016) Association between whole grain intake and all-cause mortality: a meta-analysis of cohort studies. *Oncotarget* 7: 61996-62005.
  34. Zong G, Gao A, Hu FB, Sun Q (2016) Whole Grain Intake and Mortality From All Causes, Cardiovascular Disease, and Cancer: A Meta-Analysis of Prospective Cohort Studies. *Circulation* 133: 2370-2380.
  35. Li B, Zhang G, Tan M, Libo Zhao, Lei Jin, et al. (2016) Consumption of whole grains in relation to mortality from all causes, cardiovascular disease, and diabetes: Dose-response meta-analysis of prospective cohort studies. *Medicine* 95: e4229.
  36. Ma LorenaLuna-Guevara, Juan JoséLuna-Guevara, Paola Hernández-Carranza, HéctorRuíz-Espinosa, Carlos Enrique Ochoa-Velasco (2019) Phenolic Compounds: A Good Choice against Chronic Degenerative Diseases. *Studies in Natural Products Chemistry* 59: 79-108.
  37. Gani A, Wani SM, Masoodi FA, Hameed G (2012) Whole-Grain Cereal Bioactive Compounds and Their Health Benefits: A Review. *J Food Process Technol* 3: 146.
  38. Rao S, Santhakumar AB, Chinkwo KA, Blanchard CL (2018) Q-TOF LC/MS identification and UHPLC-Online ABTS antioxidant activity guided mapping of barley polyphenols. *Food Chemistry* 266: 323-328.
  39. Pasha I, Riaz A, Saeed M, Randhawa MA (2015) *Journal of Food Processing and Preservation* 39: 1089-1097.
  40. Dei Cas M, Ghidoni R (2018) Cancer prevention and therapy with polyphenols: Sphingolipid-mediated mechanisms. *Nutrients* 10: 7.
  41. Nyström L, Lampi AM, Andersson A, Kamal-Eldin A, Gebruers K, et al. (2008) Phytochemicals and dietary fiber components in rye varieties in the HEALTHGRAIN diversity screen. *J Agric Food Chem* 56: 9758-9766.
  42. Weidner S, Amarowicz R, Karamać M, Dąbrowski G (1999) Phenolic acids in caryopses of two cultivars of wheat, rye and triticale that display different resistance to pre-harvest sprouting. *Eur Food Res Tech* 210: 109-113.
  43. Heiniö RL, Liukkonen K, Myllymäki O, Pihlava JM, Adlercreutz H, et al. (2008) Quantities of phenolic compounds and their impacts on the perceived flavour attributes of rye grain. *J Cereal Sci* 47: 566-575.
  44. Smeds A, Eklund P, Sjöholm R, Willför S, Nishibe S, et al. (2007) Quantification of a broad spectrum of lignans in cereals, oilseeds, and nuts. *J Agric Food Chem* 55: 1337-1346.
  45. Hanhineva K, Rogachev I, Aura AM, Aharoni A, Poutanen K, et al. (2012) Identification of novel lignans in the whole grain rye bran by non-targeted LC-MS metabolite profiling.

- Metabolomics 8: 399-409.
46. Liukkonen KH, Katina K, Wilhelmson A, Myllymäki O, Lampi AM, et al. (2003) Process-induced changes on bioactive compounds in wholegrain rye. *Proc Nutr Soc* 62: 117-122.
  47. Glitsø LV, Mazur W, Adlercreutz H, Wähälä K, Mäkelä T, et al. (2000) Intestinal metabolism of rye lignans in pigs. *British J Nutr* 84: 429-437.
  48. Stasiuk M, Kozubek A (2010) Biological activity of phenolic lipids. *Cell Mol Life Sci* 67: 841-860.
  49. García S, García C, Heinzen H, Moyna P (1997) Chemical basis of the resistance of barley seeds to pathogenic fungi. *Phytochemistry* 44: 415-418.
  50. Dey ES, Ahmadi-Afzadib M, Nybomb H, Tahir I (2013) Alkylresorcinols isolated from rye bran by supercritical fluid of carbon dioxide and suspended in a food-grade emulsion show activity against *Penicillium expansum* on apples. *Arch Phytopathology Plant Protect* 46: 105-119.
  51. Tahir I, Ahmadi-Afzadi M, Nybom H, Dey E (2014) Rye bran alkylresorcinols inhibit growth of *Penicillium expansum* and *Neofabraea perennans* in vitro and in vivo on different apple cultivars *Eur J Horticult Sci* 79: 218-225.
  52. Liu Z, Liu Y, Pu Z, Jirui Wang, Youliang Zheng, et al. (2013) Regulation, evolution, and functionality of flavonoids in cereal crops, *Biotechnol Lett* 35: 1765.
  53. Brazier-Hicks M, Evans K, Gershter M, Puschmann H, Steel P, et al. (2009) The C-glycosylation of flavonoids in cereals. *J Biol Chem* 284: 17926-17934.
  54. Zieliński H, Ceglińska A, Michalska A (2007) Antioxidant contents and properties as quality indices of rye cultivars. *Food Chem* 104: 980-988.
  55. Michalska A, Ceglińska A, Zieliński H (2007) Bioactive compounds in rye flours with different extraction rates. *Eur Food Res Tech* 225: 545-551.
  56. Ivanišova E, Ondrejovič M, Šilhár S (2012) Antioxidant activity of milling fractions of selected cereals. *Nova Biotechnologica ET Chimica* 11: 45-56.
  57. Del Rio D, Rodriguez-Mateos A, Spencer J, Tognolini M, Borges G, et al. (2013) Dietary (poly) phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxid Redox Signal* 18: 1818-1892.
  58. Rodriguez-Mateos A, Vauzour D, Krueger C, Shanmuganayagam D, Reed J, et al. (2014) Bioavailability, bioactivity and impact on health of dietary flavonoids and related compounds: an update. *Arch Toxicol* 88: 1803-1853.
  59. Katina K, Liukkonen K, Kaukovirta-Norja A, Adlercreutz H, Heinonen S, et al. (2007) Fermentation-induced changes in the nutritional value of native or germinated rye. *J Cereal Sci* 46: 348-355.
  60. Pedersen HA, Laursen B, Mortensen A, Fomsgaard IS (2011) Bread from common cereal cultivars contains an important array of neglected bioactive benzoxazinoids. *Food Chem* 127: 1814-1820.
  61. Tanwir F, Fredholm M, Gregersen P, Fomsgaard IS (2013) Comparison of the levels of bioactive benzoxazinoids in different wheat and rye fractions and the transformation of these compounds in homemade foods. *Food Chem* 141: 444-450.
  62. Buchmann CA, Nersesyan A, Kopp B, Schauburger D, Darroudi F, et al. (2007) Dihydroxy-7-methoxy-1, 4-benzoxazin-3-one (DIMBOA) and 2, 4-dihydroxy-1, 4-benzoxazin-3-one Gramineae are potent aneugens in human-derived liver cells (HepG2). *Cancer Lett* 246: 290-299.
  63. Adhikari KB, Tanwir F, Gregersen P, Steffensen S, Jensen B, et al. (2015) Benzoxazinoids: Cereal phytochemicals with putative therapeutic and health-protecting properties. *Mol Nutr Food Res* 59: 1324-1338.
  64. Steffensen S, Pedersen H, Adhikari KB, Laursen B, Jensen C, et al. (2016) Benzoxazinoids in prostate cancer patients after a rye-intensive diet: methods and initial results. *J Agric Food Chem* 64: 8235-8245.
  65. Peterson D, Emmons C, Hibbs A (2001) Phenolic antioxidants and antioxidant activity in pearling fractions of oat groats. *J Cereal Sci* 33: 97-103.
  66. Ndolo VU, Beta T (2013) Distribution of carotenoids in endosperm, germ, and aleurone fractions of cereal grain kernels. *Food Chem* 139: 663-671.
  67. El-Sayed M, Abdel-Aal J, Christopher Young, Iwona Rabalski, Pierre Hucl, et al. (2007) Identification and Quantification of Seed Carotenoids in Selected Wheat Species. *J Agric Food Chem* 55: 787-794.
  68. Abdoulaye C, Brou K, Jie C (2011) Phytic Acid in Cereal Grains: Structure, Healthy or Harmful Ways to Reduce Phytic Acid in Cereal Grains and Their Effects on Nutritional Quality. *American J Plant Nutrition and Fertilization Tech* 1: 1-22.
  69. Jones PJH, Ntanos FY, Raeini-Sarjaz M, Vanstone CA (1999) Cholesterol-Lowering Efficacy of a Sitostanol-Containing Phytosterol Mixture with a Prudent Diet in Hyperlipidemic Men. *Am J Clin Nutr* 69: 1140-1150.
  70. Sierksma A, Weststrate JA, Meijer GW (1999) Spreads Enriched with Plant Sterols, Either Esterified 4, 4-Dimethylsterols or Free 4-Desmethylsterols, and Plasma Total- and LDL-Cholesterol Concentrations. *Br J Nutr* 82: 273-282.
  71. Moghadasian MH, Frohlich JJ (1999) Effects of Dietary Phytosterols on Cholesterol Metabolism and Atherosclerosis:

- Clinical and Experimental Evidence. *Am J Med* 107: 588-594.
72. Law MR (2000) Plant Sterol and Stanol Margarines and Health. *West. J. Med.* 173: 43–47
73. Moreau RA, Norton RA, Hicks KB (1999) Phytosterols and Phytostanols Lower Cholesterol. *Inform* 10: 572-577.
74. Jiang Y, Wang T (2005) Phytosterols in cereal by-products. *J Amer Oil Chem Soc* 82: 439.
75. Tiwari U, Cummins E (2009) Nutritional importance and effect of processing on tocopherols in cereals. *Trends in Food Science & Technology* 20: 511-520.
76. Nielson MM, Hansen A (2008) Rapid high-performance liquid chromatography determination of tocopherols and tocotrienols in cereals. *Cereal Chemistry* 85: 248-251.
77. Finocchiaro F, Ferrari B, Gianinetti A, Dall'Asta C, Galaverna G, et al. (2007) Characterization of antioxidant compounds of red and white rice and changes in total antioxidant capacity during processing. *Molecular Nutrition & Food Research* 51: 1006-1019.
78. Panfili G, Fratianni A, Irano M (2003) Normal phase high performance liquid chromatography method for the determination of tocopherols and tocotrienols in cereals. *Journal of Agricultural and Food Chemistry* 51: 3940-3944.
79. Ryynänen M, Lampi AM, Salo-Väänänen P, Ollilainen V, Piironen V (2004) A small-scale sample preparation method with HPLC analysis for determination of tocopherols and tocotrienols in cereals. *Journal of Food Composition and Analysis* 17: 749-765.
80. Guleria P, Kumari S, Dangi N (2015)  $\beta$ -glucan: Health Benefits and Role in Food Industry - A Review. *International Journal of Enhanced Research in Science, Technology & Engineering* 4: 255-263.
81. Johansson L, Tuomainen P, Ylinen M, Ekholm P, Virkki L (2004) Structural analysis of watersoluble and -insoluble  $\beta$ -D-glucans of whole-grain oats and barley. *Carbohydrate Polymers* 58: 267-274.
82. Wood PJ, Fulcher RG, Stone BA (1993) Studies on the specificity of interaction of cereals cell wall components with congo red and calcofluor: Specific detection and histochemistry of (1 $\rightarrow$ 3),(1 $\rightarrow$ 4)-  $\beta$ -D-glucan. *Journal of Cereal Science* 1: 95-110.
83. Michaela Havrlentová, Ján Kraic (2006) Content of  $\beta$ -D-glucan in cereal grains. *Journal of Food and Nutrition Research* 45: 97-103.
84. Pomeroy S, Tupper R, Cehun-Aders M, Nestel P (2001) Oat  $\beta$ -glucan lowers total and LDL-cholesterol. *Australian Journal of Nutrition and Dietetics* 58: 51-55.
85. Laskowski W, Górska-Warsewicz H, Rejman K, Czczotko M, Zwolińska J (2019) How Important are Cereals and Cereal Products in the Average Polish Diet? *Nutrients* 11: 679.
86. Shotwell MA, Afonso C, Davies E, Chesnut RS, Larkins BA (1988) Molecular characterization of oat seed globulins. *Plant Physiol* 87: 698-704.
87. Shewry PR, Halford NG (2002) Cereal seed storage proteins: structures, properties and role in grain utilization, *Journal of Experimental Botany*. 53: 947–958.
88. Wrigley CW (2010) Cereal - grain morphology and composition In: Wrigley CW, Batey IL (Eds.). *Cereal Grains Assessing and Managing Quality*. Woodhead Publishing Series in Food Science, Technology and Nutrition. Cambridge, UK.
89. Shotwell MA (1999) Oat globulins In: Shewry, PR., Casey R (Eds.) *Seed Proteins*. Kluwer academic publishers, Netherlands.
90. Miller G, Prakash A, Decker E (2002) Whole-grain micronutrients. In Marquart, L, Slavin, JL and Fulcher, RG (Eds.) *Whole-Grain Foods in Health and Disease* St Paul MN: Eagan Press.
91. Keenan JM, Goulson M, Shamliyan T, Knutson N, Kolberg L, et al. (2007) The effects of concentrated barley beta-glucan on blood lipids in a population of hypercholesterolaemic men and women. *Br J Nutr* 97: 1162-1168.
92. Jacobs DR Jr, Andersen LF, Blomhoff R (2007) Whole-grain consumption is associated with a reduced risk of noncardiovascular, noncancer death attributed to inflammatory diseases in the Iowa Women's Health Study. *Am J Clin Nutr* 85: 1606-1614.
93. Li L, Shewry PR, Ward JL (2008) Phenolic acids in wheat varieties in the heath grain diversity screen, *Journal of Agricultural food chemistry* 56: 9732-9739.
94. Jacela JY, JM DeRouchey, MD Tokach, RD Goodband, JL Nelssen, (2010) Feed additives for swine: Fact sheets - prebiotics and probiotics and phytochemicals. *J Swine Health Prod* 18: 87-91.
95. Zhou RJ, JW Jr Erdman (1995) Phytic acid in health and disease. *Crit Rev Food Sci Nutr* 35: 495-508.
96. Saini A, Panwar D, Panesar PS and Bera MB. Bioactive Compounds from Cereal and Pulse Processing Byproducts and Their Potential Health Benefits. *Austin J Nutr Metab.* 2019; 6(2): 1068.

**Copyright:** ©2020 Remya Mohanraj. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.