

Severe Acneiform Facial Eruption: An Updated Prevention, Pathogenesis and Management

Chan Kam Tim Michael

Private Dermatologist, Hong Kong SAR.

Corresponding author

Dr. Chan Kam Tim Michael, Private Dermatologist, Hong Kong SAR. E-mail: pioneerskin@gmail.com.

Submitted: 11 June 2017; Accepted: 19 June 2017; Published: 30 June 2017

Abstract

Acne scarrings and papulopustular rosacea (PPR) are well documented cutaneous condition associated with major psychosocial morbidity. The burden of disease to the family and society is significant. A positive family history is a predictor. Inflammation involved an interplay of body innate immunity and pro-inflammatory mediators, cytokines, neuropeptides and defence immune response to microbiomes results acneiform eruption. Modern research in molecular biology, neuroimmunology and clinical science enable the practicing physician to understand more about the pathogenesis of this complex skin disease and hence better therapeutic measures and management of the disease.

Introduction

Acneiform facial eruption is a common and complex skin disease which primarily involve the epidermis and pilosebaceous units with significant co-morbidities of other body systems. In the past, clinicians have mainly focused on pathogenic microbes like *Propionibacterium acnes* by using antibiotics in treating acne vulgaris patients. The sole use of systemic and topical antibiotics has added velocity to the worldwide emergence of antibiotics-resistance skin bacteria [1-3]. Recent advances in neurophysiology, immune endocrinology and molecular biology provide new insights that inflammation through its mediators is the key element in this disease [4-6]. The following update will discuss these recent scientific findings and hopefully enable the clinicians a new mind frame in approaching the management of this chronic distressing condition. The article will mainly discuss acne vulgaris, rosacea and drug induced acneiform eruption.

Acne vulgaris is one of the earliest sign of puberty and comedone is the characteristic sign [7]. The exact cause of comedone is unknown. Proliferation of the basal layer of the epidermis and hyperkeratinisation of the acroinfundubulum of the sebaceous glands are the initial triggering factors. The excess keratin with sebum production will then cause skin pore inflammation and blockage. Overgrowth of the normal bacterial flora including *P. acnes* with secondary colonization of *Staphylococcus epidermidis* and *Staphylococcus aureus* attack the body innate and immune defence mechanism resulting in a plethora of acne manifestations like inflammatory comedones, papules, pustules, nodules, cysts, complications like hypertrophic and keloid scars. Results can be quite serious to a teenager without proper management of the disease. Acne in many studies can cause psychiatric morbidities like depression (30%), anxiety disorders (51%), low self-esteem (66%) and rarely suicidal ideation [8-10]. Acne scarring has been documented to significantly affect one's life and career [11]. Acne management need to address both the primary skin condition and

psychiatric manifestations. But without knowing details in the underlying pathogenesis of acne vulgaris; it would be difficult to enable the clinicians to counsel their patients and manage the disease.

Rosacea characterised by flushing, centrefacial erythema of face, burning and stinging sensation aggravated by heat, hot food, spices, alcohol, solar exposure and stress share a similar complexity with acne vulgaris. Rosacea sufferers are more seen in their third and fourth decade of life and the female gender [12]. An article published in the Lancet interactive journal studied Rembrandt using his self-portrait drawn in 1645 at the age of 65 arrived the conclusion that the famous Dutch painter may suffer from rosacea and other systemic diseases [13,14]. Although the picture has not given us vivid descriptions of different subtypes of rosacea: erythematous, papulopustular, phymatous, ocular and peri-oral; and the signs of cardiovascular disease in this picture are disputable; it impressed the audiences an unhappy, depressed and helpless life of the artist before his death (Figure 1) [13,14].



Figure 1: An article published in the Lancet Interactive journal studied Rembrandt using his self-portraits and came to the conclusion that Rembrandt may had had rosacea.

use of retinoids in controlling severe acne and even curing it [30]. Retinoids has been shown to interact with RXR receptors and RAR receptors which are closely associated with PPAR alpha, beta and gamma inside the nucleus of the cell. Retinoids thus exert its effects on severe acne by modulating and normalizing skin proliferation, differentiation and controlling inflammation by influencing fatty acid and lipid homeostasis [21,32]. In recent years, more and more anti-inflammatory agents have been shown to be efficacious in controlling severe acne. For example, Nicotinamide (Vitamin B3) an anti – inflammatory agent and laser delivered gold nano - particles and have been shown to improve acne outcome [33,34].

On the other hand, in Rosacea, there appears a more complicated interaction between the brain, nervous system, cardiovascular system, the gut and the skin[35,36]. A neurovascular coupling appears to occur in the pathogenesis of Rosacea. (Figure 3) Known triggering factors of rosacea refer as stressors like species, stress, exercises and heat act through the Transient Receptor Potential Vanilloid -1 (TRPV) and Transient Receptor Potential Ankyrin (TRPA) located on neuronal endings sensitize the autonomic nervous system both the sympathetic and parasympathetic pathways causing vasodilatation like flushing and hyperalgesia like stinging sensation [37,38]. Nerves containing Calcitonin genes related peptide (CGRP) may be linked to flushing through the messenger cAMP. Furthermore, CGRP may also serve to regulate inflammation by modulating chemokine production by the cells lining the blood vessels. TRPV act like a N-methyl-D-aspartate (NMDA) receptor cause calcium influx thus changes transmembrane potential transmits signals to the cardiovascular system via the central nervous system. Excessive solar radiation generates Fibroblast Growth Factor (FGF) and Vascular Epidermal Growth Factor (VEGF) promoting blood vessel proliferation

and skin and pilosebaceous gland inflammation resulting in skin papules, pustules. The inflammatory insults also cause a direct vaso - vascular effects on angiogenesis resulting in erythema and telangiectasia respectively [39,40]. Pathogenic microorganisms may invade the defective epidermal barrier of the skin of rosacea and the epidermal keratinocyte in response produce a potent anti-bacterial cytokine called beta defensins [41]. Cathelicidin (LL-37); a potent antibacterial protein by skin keratinocyte in response to antimicrobial invasion; is produced through the Kallikrein 5 pathway involving the Matrix metalloproteinases (MMP). LL-37 also known as defensins activated protein has been demonstrated to be a major pathogenetic factor in causing rosacea through inflammation and angiogenesis of skin [42-45]. Finally, an abnormal skin microflora consisting of an increasing number of Demodex folliculorum in the sebaceous glands and S. epidermidis in the defective epidermal barrier initiate the production of Toll –Like Receptor (TLR) from keratinocytes which also enhance the production of LL-37 [46,47]. Recently, there are study suggests that the bacterium Bacillus oleronius harboured by the Demodex mites are the genuine culprit in causing the release of TLR [48]. Whether the Demodex play a more significant role in the pathogenesis of rosacea than LL-37 is not known; however; this may be important especially when one is considering using an antimicrobial to treat rosacea. Nonetheless, LL-37 and TRPV/ TRPA channels are important neurovascular mediators and signalling pathways in causing rosacea. This may have important implications in designing future pharmacological therapy in managing severe rosacea as LL-37 inhibitors and TRPV channels blockers may be found to antagonize the neurovascular coupling that occur in rosacea [49].




Figure 3: Pathways known to contribute to the pathophysiology and clinical manifestations of Rosacea (From Two AM, Wu W, Gallo RL, Hata TR. Rosacea: part I. Introduction, categorization, histology, pathogenesis, and risk factors. JAAD.2015;72:749-58).

Many drugs have been reported to adversely affected the treated patients with severe acneiform skin eruption [50]. They can be classified as hormonal, anti-depressant, anti-convulsant and anti-cancer drugs. They may result a severe papulopustular skin eruption resembling acne and rosacea. The pathogenesis may also involve an aberrant immune endocrine metabolic signalling pathways in genetically susceptible individuals. Epidermal growth factor receptor (EGFR) inhibitors and tyrosine kinase inhibitors which are widely used biologics targeting specific receptors of the cancer cells have all reported to cause significant skin reactions especially acneiform eruptions (Table 1) [51-54].

Table 1: Drugs that are known to induce Acneiform facial eruption

	Drugs that caused acneiform facial eruption
Steroid Hormones	Topical steroids
	Systemic steroids
	Anabolic steroids for muscle building
	Testosterone
Antidepressants	Lithium
Antiepileptic	Phenytoin
Halogens	Iodides
	Bromides
Target therapy for metastatic internal malignancies	Anti EGFR
	Anti BRAF

The recent development in the research of molecular biology reveal a better understanding of the pathogenesis of severe acneiform eruption compatible with clinical observations and findings. This may assist the busy clinicians battling with this distressing disease to have a new, more positive, dualistic mind frame in approaching the management. The conventional use of antibiotic alone in treating severe cases of these patients may not be all satisfactory. It will only further promote the already devastating bacteria antibiotic resistance pandemic without relieving the actual psychological distress of these sufferers. Prevention is better than cure. Counselling the teens suffered from severe acne and a heavy work laden middle-aged women suffered from rosacea required an updated knowledge of what's happening in acne and rosacea [55]. An effective approach in future may involve dietary manipulation by using pro-biotics; anti-inflammatory agents; cytokines inhibitors and genuine receptors antagonist to target the multi- faceted of the pathogenesis of acneiform eruption.

In conclusion, a new mind frame need to be set to understand severe acneiform eruption. The burden of disease of acne and rosacea is more than skin deep; and may neurologically, psychologically and systemically affect an individual. It involves an orchestral interplay between the gut, immune system, vascular system neuro-endocrine and brain organs. Further advancement in understanding the pathogenesis hence management of this complex disease entity required collaboration, communication and inter connectiveness involving the neuro-psychiatrists, neuroscientists, immune endocrinologists and all relevant medical professional partners.

Acknowledgement

This article is part of the Keynote presentation in the Joint Conference of 12th International Conference on Neurology and Neurophysiology & 2nd International Conference and Exhibition on Dual Conference; May 18-20, 2017 Munich Germany.

References

- Cooper AJ (1998) Systemic review of Propionibacterium acne resistance to systemic antibiotics. *Med J Austral* 169: 259-261.
- Miller YW, Eady EA, Lacey RW, Cove JH, Joanes DN, et al. (1996) Sequential antibiotic therapy for acne promotes the carriage of resistant staphylococci on the skin of contacts. *J Antimicrob Chemother* 38: 829-837.
- Fanelli M, Kupperman E, Lautenbach E, Edelstein PH, Margolis DJ (2011) Antibiotics, Acne, and Staphylococcus aureus Colonization. *Arch Dermatol* 147: 917-921.
- Emil A. Tanghetti (2013) the Role of Inflammation in the Pathology of Acne. *J Clin Aesthet Dermatol* 6: 27-35.
- Kim J (2005) Review of the innate immune response in acne vulgaris: activation of Toll-like receptor 2 in acne triggers inflammatory cytokine responses. *Dermatology* 211: 193-198.
- Do TT, Zarkhin S, Orringer JS, Nemeth S, Hamilton T, et al. (2008) Computer-assisted alignment and tracking of acne lesions indicate that most inflammatory lesions arise from comedones and. de novo. *J Am Acad Dermatol* 58: 603-608.
- Cunliffe WJ, Holland DB, Jeremy A (2004) Comedone formation: etiology, clinical presentation, and treatment. *Clin Dermatol* 22: 367-374.
- Daderwale D (2016) Acne more than skin deep - Psychiatric and self - esteem in acne patients. *European Journal of Pharmacueticals and Medical Research* 3: 532 -538.
- Barnes LE, Levender MM, Fleischer AB Jr, Feldman SR (2012) Quality of life measures for acne patients. *Dermatol Clin* 30: 293-300.
- Halvorsen JA, Stern RS, Dalgard F, Thoresen M, Bjertness E, Lien L (2011) Suicidal ideation, mental health problems, and social impairment are increased in adolescents with acne: a population-based study. *J Invest Dermatol* 131: 363-370.
- Williams HC, Dellavalle RP, Garner S (2012) Acne vulgaris. *Lancet* 379: 361-372
- Kyriakis KP, Palamaras I, Terzoudi S, Emmanuelides S, Michailides C, et al. (2005) Epidemiologic aspects of rosacea. *JAAD* 53: 918-919.
- Duff R F, A Zoltnick, Graeme J Hanke (1998) Rembrandt's self-portrait. *The Lancet. Correspondence* 351: 915.
- Espinel CH (1997) A medical evaluation of Rembrandt. His self-portrait: ageing, disease, and the language of the skin. *Lancet* 350: 1835-1837.
- Huynh T T (2013) The Psych Impact of Rosacea. *Am Health & drug Benefits* 6: 6.
- Rainer BM, Fischer AH, Luz Felipe da Silva D, Kang S, Chien AL (2015) Rosacea is associated with chronic systemic diseases in a skin severity-dependent manner: results of a case-control study. *J Am Acad Dermatol* 73: 604-608.
- Egeberg A, Ashina M, Gaist D, Gislason GH, Thyssen JP, et al. (2017) Prevalence and risk of migraine in patients with rosacea: a population- based cohort study. *Journal of the American Academy of Dermatology* 76: 454-458.
- Spoendlin J, Karatas G, Furlano RI, Jick SS, Meier CR (2016) Rosacea in patients with ulcerative colitis and Crohn's disease: a population-based case-control study. *Inflammatory*

- Bowel Disease 22: 680-687.
19. Miri Kim , Kwang Hyun Choi , Se Won Hwang, Young Bok Lee, Hyun Jeong Park, et al. (2016) Inflammatory bowel disease is associated with an increased risk of inflammatory skin diseases: a population-based cross-sectional study. *Journal of the American Academy of Dermatology* 76: 40-48.
 20. Egeberg A, Fowler JF Jr, Gislason GH, Thyssen JP (2017) Rosacea and risk of cancer in Denmark. *Cancer Epidemiol* 47: 76-80.
 21. Egeberg A, Fowler JF Jr, Gislason GH, Thyssen JP (2016) Nationwide Assessment of Cause-Specific Mortality in Patients with Rosacea: A Cohort Study in Denmark. *Am J Clin Dermatol* 17: 673-679.
 22. Drake L (2017) National Rosacea Society Newsletter. Glioma may associate with Rosacea.
 23. Whitney P, Bowe, Alan C, Logan (2011) Acne vulgaris, probiotics and the gut-brain-skin axis – back to the future? *Gut Pathogens* 3: 1.
 24. Smith RN, Mann NJ, Braue A, Makelainen H, Varigos GA (2007) The effect of a high-protein, low glycemic-load diet versus a conventional, high glycemic-load diet on biochemical parameters associated with acne vulgaris: a randomized, investigator-masked, controlled trial. *J Am Acad Dermatol* 57: 247-256.
 25. Zouboulis CC, Böhm M (2004) Neuroendocrine regulation of sebocytes -- a pathogenetic link between stress and acne. *Exp Dermatol* 13: 31-35.
 26. Smith TM, Cong Z, Gilliland KL, Clawson GA, Thiboutot DM (2006) Insulin-like growth factor-1 induces lipid production in human SEB-1 sebocytes via sterol response element-binding protein-1. *J Invest Dermatol* 126: 1226-1232.
 27. Fabbrocini G, Izzo R, Faggiano A, Del Prete M, Donnarumma M, et al. (2016) Clin Exp Derm. Low glycaemic diet and metformin therapy: a new approach in male subjects with acne resistant to common treatments 41: 38-42.
 28. Trivedi NR, Cong Z, Nelson AM, Albert AJ, Rosamilia LL, et al. (2006) Peroxisome proliferator-activated receptors increase human sebum production. *J Invest Dermatol* 126: 2002-2009.
 29. Maryam Ahmadian, Jae Myoung Suh, Nasun Hah, Christopher Liddle, Annette R Atkins, et al. (2013) PPAR γ signaling and metabolism: the good, the bad and the future. *Nature Medicine* 99: 557-566.
 30. Zaenglein AL, Pathy AL, Schlosser BJ, Alikhan A, Baldwin HE, et al. (2016) Guidelines of care for the management of acne vulgaris. *J Am Acad Dermatol* 74: 945-973.
 31. Thielitz A, Gollnick H (2008) Topical retinoids in acne vulgaris: update on efficacy and safety. *Am J Clin Dermatol* 9: 369-381.
 32. Zaenglein AL (2008) Topical retinoids in the treatment of acne vulgaris. *Semin Cutan Med Surg* 27: 177-182.
 33. Grange PA, Raingeaud J, Calvez V, Dupin N (2009) Nicotinamide inhibits Propionibacterium acnes-induced IL-8 production in keratinocytes through the NF-kappaB and MAPK pathways. *J Dermatol Sci* 56: 106-112.
 34. Dilip Y Paithankar, Fernanda H Sakamoto, William A Farinelli, Kosiratna G, Blomgren RD, et al. (2015) Acne Treatment Based on Selective Photothermolysis of Sebaceous Follicles with Topically Delivered Light-Absorbing Gold Microparticles. *Journal of Investigative Dermatology* (2015) 135: 1727-1734.
 35. Drake L (1998) Rosacea takes emotional toll. *Rosacea Rev* summer: 2.
 36. Sobottka A; Lehmann P. Rosacea (2009) new advances in pathophysiology, clinical staging and therapeutic strategies. *Hautarzt* 60: 999-1009
 37. Two AM, Wu W, Gallo RL, Hata TR (2015) Rosacea: part I. Introduction, categorization, histology, pathogenesis, and risk factors. Pathways known to contribute to the pathophysiology and clinical manifestations of Rosacea. *JAAD* 72: 749-758.
 38. Sulk M, Seeliger S, Aubert J, Schwab VD, Cevikbas F, et al. (2011) Distribution and expression of non-neuronal transient receptor potential (TRPV) ion channels in rosacea. *Journal of Investigative Dermatology* 132: 1253-1262.
 39. Helfrich YR, Maier LE, Cui Y, Fisher GJ, Chubb H, et al. (2015) J. Clinical, histologic, and molecular analysis of differences between erythematotelangiectatic rosacea and telangiectatic photoaging. *JAMA Dermatol* 151: 825-836.
 40. Schwab VD, Sulk M, Seeliger S, Nowak P, Aubert J, et al. (2011) Neurovascular and neuroimmune aspects in the pathophysiology of rosacea. *Journal of Investigative Dermatology Symposium Proceedings* 15: 53-62.
 41. Zhou M, Xie H, Cheng L, Li J (2016) Clinical characteristics and epidermal barrier function of papulopustular rosacea: a comparison study with acne vulgaris. *Pak J Med Sci* 32: 1344-1348.
 42. Yamasaki K, DiNardo A, Bardan A, Masamoto Murakami, Takaaki Ohtake, et al. (2007) Increased serine protease activity and cathelicidin promotes skin inflammation in rosacea. *Nature Medicine* 13: 975-980.
 43. Schaubert J, Gallo RL (2008) Antimicrobial peptides and the skin immune defense system. *J Allergy Clin Immunol* 122: 261-266.
 44. Navarini AA, Meller S, Gilliet M, Conrad C (2010) Antimicrobial peptide-DNA complexes are implicated in initial pathogenesis of rosacea. *Journal of Investigative Dermatology* 130: 1-13.
 45. Reinholz M, Ruzicka T, Schaubert J (2012) Cathelicidin LL-37; an antimicrobial peptide with a role in inflammatory skin disease. *Ann Dermatol* 24: 126 -135.
 46. Murillo N, Aubert J, Raoult D (2014) Microbiota of Demodex mites from rosacea patients and controls. *Microb Pathog* 72: 37-40.
 47. Whitfeld M, Gunasingam N, Leow LJ, Shirato K, Preda V (2011) Staphylococcus epidermidis: a possible role in the pustules of rosacea. *Journal of the American Academy of Dermatology* 64: 49-52.
 48. Jarmuda S, McMahon F, Zaba R, O'Reilly N, Jakubowicz O, et al. (2014) Correlation between serum reactivity to Demodex-associated Bacillus oleronius proteins, and altered sebum levels and Demodex populations in erythematotelangiectatic rosacea patients. *J Med Microbiol* 63: 258-262.
 49. Sing D, Vaughan R, Koo (2014) CC. LL-37 peptide enhancement of signal transduction by Toll-like Receptor is regulated by pH, identification of a peptide antagonist of LL 37. *J Biol Chem* 289: 27614-27624.
 50. Momin SB, Peterson A, Del Rosso JQ (2010) A status report on drug-associated acne and acneiform eruptions. *J Drugs Dermatol* 9: 627-636.
 51. Eames T, Landthaler M, Karrer S (2007) Severe acneiform skin reaction during therapy with erlotinib (Tarceva), an epidermal growth factor receptor (EGFR) inhibitor. *Eur J Dermatol* 17: 552-553.

-
52. Journagan S, Obadiah J (2006) An acneiform eruption due to erlotinib: prognostic implications and management. *J Am Acad Dermatol* 54: 358-360.
 53. Myskowski PL, Halpern AC (2009) Skin reactions to the new biologic anticancer drugs. *Curr Opin Support Palliat Care* 3: 294-299.
 54. Jung YS, Kim M, Lee JH, Cho BK, Kim DW (2015) Acneiform eruptions caused by various second-generation tyrosine kinase inhibitors in patients with chronic myeloid leukaemia. *Br J Dermatol* 174: 456-458.
 55. Drake L (2008) National Rosacea Society Newsletter. Women May Need Added Therapy.

Copyright: ©2017 Dr. Chan Kam Tim Michael. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.